


Chemistry 344: Spectroscopy and Spectrometry Problem Set 1

Name (print): _____


TA Name (print): _____

most important
↓
region for Chem344

- I. The IR spectrum below is for isopropanol; identify any useful IR absorptions $>1500\text{ cm}^{-1}$. Based upon the spectrum, comment on whether you expect that it was a pure sample or diluted in a different solvent.




- II. Using the mass spectrum of 1-propanol shown below, answer the questions that follow about its fragmentation. Remember that you are not expected to interpret all signals in a mass spectrum.




- A. Why does the peak at $m/z = 15$ have such a low relative intensity?
- B. Draw a valid resonance structure of the molecular ion ($m/z = 60$). Draw a molecular ion responsible for the small peak at $m/z = 61$.

- C. In fragmentations of cations and radical cations, a single molecular (parent) ion can give rise to several fragments. Provide fragmentation mechanisms from the molecular ion ($m/z = 60$) that will rationalize the formation of ions with values of $m/z = 59$, 42, and 31. For clarity, it may help to show each fragmentation in a different color




- D. What can you conclude about the formation and decomposition of the ion responsible for the signal at $m/z = 31$?

- III. Using the mass spectrum of 2,2,4-trimethylpentane (isooctane, C_8H_{18}) shown below, answer the questions that follow about its fragmentation.




SDBSWeb : <http://sdb.srioddb.aist.go.jp> (National Institute of Advanced Industrial Science and Technology)

- A. In this case, the molecular ion is of such a low intensity it is not detected. Draw the molecular ion of 2,2,4-trimethylpentane and determine its m/z value. What does the low intensity indicate about its stability?
- B. Draw a fragmentation mechanism that explains how the molecular ion can produce ions with values of $m/z = 57$ and 56.




- C. The ion responsible for the signal at $m/z = 56$ can be represented by a set of resonance structures. Depict the most important resonance structures and a resonance hybrid.


- D. An alternate fragmentation mechanism from the molecular ion can be used to rationalize the production of ions with values of $m/z = 43$, 42, and 41. Provide an electron pushing mechanism for this fragmentation pathway.



- E. Two separate fragmentations of the molecular ion can lead to ions with m/z values of 99; provide them below.





- IV. Using the mass spectrum of 2-pentanone ($C_5H_{10}O$) shown below, answer the questions that follow about its fragmentation.




- A. Draw a valid resonance structure of the molecular ion ($m/z = 86$). Explain the source of the small peak at $m/z = 87$.

- B. Important decomposition pathways for the molecular ion of carbonyl-containing compounds such as 2-pentanone involve α -cleavage. Beginning with both viable α -cleavage decomposition pathways, draw fragmentation mechanisms that lead to ions with m/z values of 71, 43, 15. Make sure your mechanisms account for two different ions with m/z values of 43.
- C. The other major decomposition pathway arises from a McLafferty rearrangement of the molecular ion followed by the loss of an ethylene gas molecule. Show an electron-pushing mechanism for this decomposition which will rationalize the peak with a value of $m/z = 58$.




- V. A mixture of regioisomers *o*-vanillin and vanillin ($C_8H_8O_3$) were analyzed by IR and GC-MS. Analyze the spectra below and answer the accompanying questions.



- A. Identify any useful IR absorptions that can help identify this sample as a mixture of *o*-vanillin and vanillin. Is it possible to use the IR of the mixture to identify each component?




- B. Use the GC-Mass spectrum of the mixture provided below to determine the ratio of vanillin to *o*-vanillin. The most intense signals of vanillin are m/z values of 152 and 106, while the most intense signals of *o*-vanillin are m/z values of 152 and 151.



- C. Provide an electron-pushing mechanism for the fragmentation of *o*-vanillin that can rationalize the presence of ions with m/z values of 152, 151, and 123. There are many other fragmentations possible, you only need to analyze the pathway that produces the indicated signals above.

- VI. Use the GC-Mass spectrum of the student product obtained from the Friedel-Crafts acylation of bromobenzene (shown on page 12) to answer the following questions. **Only signals with intensity greater than 15% relative intensity to the base peak are shown for clarity.**



- A. Which of the peaks (1 – 3) detected in the GC are the reactant(s)? Which of the peak(s) in the GC are the product(s)?
- B. In any of the mass spectra provided for GC peaks 1 – 3, how can you tell which ions contain bromine?
- C. Provide a fragmentation mechanism that will account for the signals listed below for each molecule detected in the GC trace. Identify the species most likely responsible for the following m/z signals:


Peak #1 – 158, 156, 77, 51, 50

Peak #2 – 200, 198, 185, 183, 157, 155, 76, 43

Peak #3 – 200, 198, 185, 183, 157, 155, 76, 43


D. What is the conversion percentage from reactant to the total products by GC?

E. What is the ratio of the major to minor product by GC?




Peak#	R. Time	Area	Area%
1	2.488	842786	14.73
2	6.345	143248	2.50
3	8.081	4736553	82.77
		5722587	100.00

Peak#:1 R. Time:2.5(Scan#:30) MassPeaks:



Peak#:2 R. Time:6.3(Scan#:492) MassPeaks:13



Peak#:3 R. Time:8.1(Scan#:701) MassPeaks:1

