


**Chemistry 344: Spectroscopy and Spectrometry Problem Set 1**


Name (print): \_\_\_\_\_

TA Name (print): \_\_\_\_\_

- I. The IR spectrum below is for isopropanol; identify any useful IR absorptions  $>1500\text{ cm}^{-1}$ . Based upon the spectrum, comment on whether you expect that it was a pure sample or diluted in a different solvent.




- II. Using the mass spectrum of 1-propanol shown below, answer the questions that follow about its fragmentation. Remember that you are not expected to interpret all signals in a mass spectrum.




- A. Why does the peak at  $m/z = 15$  have such a low relative intensity?
- B. Draw a valid resonance structure of the molecular ion ( $m/z = 60$ ). Draw a molecular ion responsible for the small peak at  $m/z = 61$ .

- C. In fragmentations of cations and radical cations, a single molecular (parent) ion can give rise to several fragments. Provide fragmentation mechanisms from the molecular ion ( $m/z = 60$ ) that will rationalize the formation of ions with values of  $m/z = 59$ , 42, and 31. For clarity, it may help to show each fragmentation in a different color




- D. What can you conclude about the formation and decomposition of the ion responsible for the signal at  $m/z = 31$ ?

- III. Using the mass spectrum of 2,2,4-trimethylpentane (isooctane,  $C_8H_{18}$ ) shown below, answer the questions that follow about its fragmentation.




SDBSWeb : <http://sdb.srioddb.aist.go.jp> (National Institute of Advanced Industrial Science and Technology)

- A. In this case, the molecular ion is of such a low intensity it is not detected. Draw the molecular ion of 2,2,4-trimethylpentane and determine its  $m/z$  value. What does the low intensity indicate about its stability?
- B. Draw a fragmentation mechanism that explains how the molecular ion can produce ions with values of  $m/z = 57$  and 56.




- C. The ion responsible for the signal at  $m/z = 56$  can be represented by a set of resonance structures. Depict the most important resonance structures and a resonance hybrid.


- D. An alternate fragmentation mechanism from the molecular ion can be used to rationalize the production of ions with values of  $m/z = 43$ , 42, and 41. Provide an electron pushing mechanism for this fragmentation pathway.



- E. Two separate fragmentations of the molecular ion can lead to ions with  $m/z$  values of 99; provide them below.





- IV. Using the mass spectrum of 2-pentanone ( $C_5H_{10}O$ ) shown below, answer the questions that follow about its fragmentation.




- A. Draw a valid resonance structure of the molecular ion ( $m/z = 86$ ). Explain the source of the small peak at  $m/z = 87$ .

- B. Important decomposition pathways for the molecular ion of carbonyl-containing compounds such as 2-pentanone involve  $\alpha$ -cleavage. Beginning with both viable  $\alpha$ -cleavage decomposition pathways, draw fragmentation mechanisms that lead to ions with  $m/z$  values of 71, 43, 15. Make sure your mechanisms account for two different ions with  $m/z$  values of 43.
- C. The other major decomposition pathway arises from a McLafferty rearrangement of the molecular ion followed by the loss of an ethylene gas molecule. Show an electron-pushing mechanism for this decomposition which will rationalize the peak with a value of  $m/z = 58$ .

- V. A mixture of regioisomers *o*-vanillin and vanillin ( $C_8H_8O_3$ ) were analyzed by IR and GC-MS. Analyze the spectra below and answer the accompanying questions.






- A. Identify any useful IR absorptions that can help identify this sample as a mixture of *o*-vanillin and vanillin. Is it possible to use the IR of the mixture to identify each component?






- B. Use the GC-Mass spectrum of the mixture provided below to determine the ratio of vanillin to *o*-vanillin. The most intense signals of vanillin are  $m/z$  values of 152 and 106, while the most intense signals of *o*-vanillin are  $m/z$  values of 152 and 151.



- C. Provide an electron-pushing mechanism for the fragmentation of *o*-vanillin that can rationalize the presence of ions with  $m/z$  values of 152, 151, and 123. There are many other fragmentations possible, you only need to analyze the pathway that produces the indicated signals above.

- VI. Use the GC-Mass spectrum of the student product obtained from the Friedel-Crafts acylation of bromobenzene (shown on page 12) to answer the following questions. **Only signals with intensity greater than 15% relative intensity to the base peak are shown for clarity.**



- A. Which of the peaks (1 – 3) detected in the GC are the reactant(s)? Which of the peak(s) in the GC are the product(s)?
- B. In any of the mass spectra provided for GC peaks 1 – 3, how can you tell which ions contain bromine?
- C. Provide a fragmentation mechanism that will account for the signals listed below for each molecule detected in the GC trace. Identify the species most likely responsible for the following  $m/z$  signals:


Peak #1 – 158, 156, 77, 51, 50

Peak #2 – 200, 198, 185, 183, 157, 155, 76, 43

Peak #3 – 200, 198, 185, 183, 157, 155, 76, 43


D. What is the conversion percentage from reactant to the total products by GC?

E. What is the ratio of the major to minor product by GC?




Peak#	R. Time	Area	Area%
1	2.488	842786	14.73
2	6.345	143248	2.50
3	8.081	4736553	82.77
		5722587	100.00

Peak#:1 R. Time:2.5(Scan#:30) MassPeaks:



Peak#:2 R. Time:6.3(Scan#:492) MassPeaks:13



Peak#:3 R. Time:8.1(Scan#:701) MassPeaks:1

