Hour Exam #2 Chemistry 343	Last Name
Professor Gellman	
2 November 2016	First Name

**General Instructions:** 

- (i) Use scratch paper at back of exam to work out answers; final answers must be recorded at the proper place on the exam itself for credit.
- (ii) Print your name on each page.
- (iii) Please keep your paper covered and your eyes on your own work.

  Misconduct will lead to failure in the course.
- 1. (13 points) Shown below are the structures of two drug molecules. In each case, CIRCLE each of the chiral centers, and indicate configuration (R vs. S) for each chiral center. Be sure to answer the question at the bottom.

(c) For each structure below, one of the three statements about optical activity given is correct. In each case, write the numeral corresponding to the correct statement in the square.

$$H_2N$$
 $H_2N$ 
 $H_2N$ 
 $H_3N$ 
 $H_4N$ 
 $H_4N$ 

- (1) The specific rotation will be identical to that of the molecule shown in part (b).
- (2) The specific rotation will have the same absolute value but the opposite sign relative to that of the molecule shown in part (b).
- (3) No prediction about the specific rotation can be made.

2. (24 points) Show the major product(s) expected from each reaction below.

(single enantiomer)

3. (12 points) Show the reagents required to convert the starting molecule to the indicated product. If necessary, differentiate between or among distinct steps by using "1)", "2)", etc.

4. (22 points)

(a) Two products are formed in the reaction indicated below. Draw the structures of those products after the arrow.

(single enantiomer)
$$F_3C^{NN}$$

$$Pd/C$$

(b) On the line below, indicate the relationship between the two product molecules. Be as specific as possible.

(c) Draw both chair conformations of each of the product molecules you indicated in part (a). For each molecule, CIRCLE the conformation you expect to be more stable.

(d) Of the products generated in the reaction above, which molecule should be most stable (i.e., which should release least heat upon combustion)? Answer this question by redrawing below the appropriate molecule.

5. (10 points) Consider the Newman projection below.

$$\begin{array}{c|c} CI & \stackrel{F}{ \longrightarrow} & \stackrel{H_2}{ \longrightarrow} & \stackrel{H}{ \longrightarrow} & \stackrel{H}{$$

(a) Provide a "hexagon" drawing for the molecule shown in the Newman projection above.

(b) Provide a "chair" drawing for the molecule shown in the Newman projection above, maintaining the conformation indicated by this Newman projection.

(c) Shown below are the Newman projections for the two chair forms of a substituted cyclohexane derivative. CIRCLE the Newman projection that corresponds to the more stable chair conformation.


$$\begin{array}{c} H \\ H \\ C \\ H \\ C \\ H_2 \\ H \end{array} \begin{array}{c} H \\ O \\ C \\ H_3 \\ \end{array}$$

6. (8 points) Provide a mechanism ("curved arrows") for the reaction shown below.


Name	

7. (11 points) Shown below are a reaction and the energy diagram for this reaction. (Note that there is only one step in this reaction; no follow-up treatment with  $(CH_3)_2S$  or  $H_2O_2$ .) Draw the structures for molecules A, B and C in the appropriate boxes below.





reaction progress



Problem #	Score
1	/13
2	/24
3	/12
4	/22
5	/10
6	/ 8
7	/11

Total:

/100

УША	2 <b>He</b> 4.00	10 Ne 20.18	18 Ar 39.95	36 <b>Kr</b> 83.80	54 <b>Xe</b> 131.29	86 Rn 222	]	
e i		9 F	17 CI 35.45	35 <b>Br</b> 79.90	53 I 126.90	85 <b>At</b> 210		
	VIA VIIA	8 <b>O</b> 16.00	16 <b>S</b> 32.07	% <b>Se</b> 78.96	$ \begin{array}{c ccccccccccccccccccccccccccccccccccc$	Po		
	VA	<b>Z</b> 10.41	15 <b>P</b> 30.97	33 <b>AS</b> 74.92	51 S <b>b</b> 121.76	83 <b>Bi</b> 208.98		
	IIIA IVA	6 C 12.01	<b>Si</b> 28.09	32 <b>Ge</b> 72.61	Sh 118.71	82 <b>Pb</b> 207.2		
	ША	5 <b>B</b> 10.81	13 <b>AI</b> 26.98	31 <b>Ga</b> 69.72	49 <b>In</b> 114.82	80 81 8 <b>Hg</b> TI 200.59 204.38		
ents			B	30 <b>Zn</b> 65.39	4	80 <b>Hg</b> 200.59	112	
Periodic Table of the Elements			æ	29 Cu 63.55	45 46 47 <b>Rh Pd Ag</b> 102.91 106.42 107.87	79 <b>Au</b> 196.97	272	
the ]	ison			28 Ni 58.69	46 <b>Pd</b> 106.42	78 Pt 195.08	110	
ole of	Mad		VIIIB	Z7 Co 58.93		77 <b>Lr</b> 192.22	109 <b>Mt</b> 266	
c Tak	UW-Madison			26 <b>Fe</b> 55.85	44 <b>Ru</b> 101.07	76 <b>Os</b> 190.23	108 Hs 265	
riodi		377	VIIB	25 Mn 54.94	$egin{array}{c} 43 \ \mathbf{Tc} \ 98 \ \end{array}$	75 <b>Re</b> 186.21	107 <b>Bh</b> 262	
Pe			VIB	24 Cr 52.00	42 <b>Mo</b> 95.94	74 <b>W</b> 183.85	106 2 <b>63</b> 263	
			VB	23 V 50.94	41 <b>Nb</b> 92.91	$\mathbf{T}^3$ 180.95	105 <b>Db</b> 262	
			IVB	22 Tri 47.88	40 <b>Zr</b> 91.22	72 <b>Hf</b> 178.49	104 <b>Rf</b> 261	8
			E E	21 <b>Sc</b> 44.96	39 Y 88.90	<i>57</i> <b>La</b> * 138.91	89 <b>Ac</b> * 227	
	ПА	4 Be 9.01	12 <b>Mg</b> 24.30	% <b>Ca</b> 40.08	38 <b>Sr</b> 87.62	56 <b>Ba</b> 137.33	88 <b>Ra</b> 226	
IA	10.1	3 <b>Li</b> 6.94	11 Na 22.99	19 <b>K</b> 39.10	<b>3.b</b> 5.47	55 Cs 132.91	87 Fr 223	

[		7	Ī		
77	ra La	174.9		7	
70	χp	173.04	102	Z	259
eg E	Tm	168.93	101	Md	258
, F	E	167.26	100	Fm	257
72		164.93	83	Es	252
88 F	C S	162.50	88	Cf	251
ිසි [	<b>-</b> }	158,	26	Bk	247
₽ 7	ָל בּי	157.25	86	Cm	247
ස දූ	מַן בּיַרְ	151.96	88	Am	243
25 D	DIE I	150.36	な	Pu	244
1		145	83	ND	237
8	מון דער	144.24	83	þ	238.03
ි සි	7.4	140.91	91	Th Pa U	231.04
58 59 60 6.	2 S	140.12	83	Th	232.04

\* Lanthanides

\*\* Actinides