Hour Exam #1
Chemistry 343
<b>Professor Gellman</b>
7 October 2013

First Name			 

## General Instructions:

(i) Use scratch paper at back of exam to work out answers; final answers must be recorded at the proper place on the exam itself for credit. Models are allowed.

Last Name

- (ii) Print your name on each page.
- (iii) Please keep your paper covered and your eyes on your own work. Misconduct will lead to failure in the course.
- 1. (9 points) For each alkene below, circle the proper stereoisomer designation.

E Z Neither

E Z Neither

E Z Neither

Name

2. (8 points) The molecule below, efavirenz, is a drug used for treatment of AIDS.

Put a CIRCLE around each sp<sup>3</sup>-hydridized carbon atom.

Put a TRIANGLE around each sp-hydridized carbon atom.

Put a SQUARE around each sp²-hydridized carbon atom that is not bonded to another carbon.

- 3. (20 points)
- (a) Draw in the other side of the Brønsted acid-base equilibrium involving the two species shown below.

(b) Draw the two bases from the equilibrium above in the space below, and CIRCLE the one that is stronger.

(c) Which side of the the equation in part (a) is favored (circle the correct word below)?

(d) Draw in the other side of the Brønsted acid-base equilibrium involving the single species below.


Name			

4. (22 points)

(a) Draw all three staggered Newman projections about the central bond of molecule A, which is shown below.

Br A

(b) Draw all three staggered Newman projections about the central bond of molecule B, which is shown below.



(c) The heats of formation for isomers A and B are negative, but the value for A is slightly more negative than the value for B (i.e., the absolute value of the heat of formation of A is larger than the absolute value of the heat of formation of B).

A bromine atom is nearly the same size as a methyl group.

Based on this information and the answers to parts (a) and (b), explain in <u>one sentence</u> why the heat of formation is more negative for A than for B.

Name		

5. (16 points) The reaction shown below starts with a single alkene, but two products are produced in comparable quantities. Provide a mechanism ("curved arrows") for this reaction. Show every atom in each structure you draw.

Name \_\_\_\_\_00225

 Problem #
 Score

 1
 / 9

 2
 / 8

 3
 / 20

 4
 / 22

 5
 / 16

 6
 / 25

\_\_\_\_

Total: /100

## Periodic Table of the Elements

			<b>Fir</b> (223)	C.S 132.91	85.47 55	37 39.10	Na 22.99	6.94	
	Lanthanides		<b>Ra</b> 226.03	137.34 88	87.62	40.08	Mg 24.31	<b>B</b> e 9.01	4
	<u></u>	7	Ac (227)	La 138.91	<b>Y</b> 88.91	Sc 44.96	21		
3	58 Ce 140.12		Unq*	178.49	91.22 72	47.90	22		H -1.008
2	\$9 <b>P1</b> : 140.91				<del>-</del>				φα
	60 Nd 144.24		Unp* (262)	Ta 180.95					
-		-	Unh*	183.85	Mo 95.94	52.00 42	24		
	6) <b>P<sub>111</sub></b> (145)		Uns*			Į.	25		
	62 <b>Sm</b> 150.35		* Uno*		<del>-  </del>				
	63 IEu 151.96			<del> </del>		<u> </u>			
1	· · · · · · · · · · · · · · · · · · ·		(266)	-			_		
	64 <b>Gd</b> 157.25			<b>Pt</b>	<b>Pd</b> 106.4	Ni 58.71			
	158.93			Au 196.97	<b>Ag</b> 107.87	Cu 63.55		<b>%</b>	
	Dy 162.50			80 Hg 200.59	Cd	30 <b>Zn</b> 65.37			
	67 <b>Ho</b> 164.93			81 <b>T1</b> 204.37	ln 114.82	31 Ga 69.72	13 A1 26.98	5 <b>B</b>	
	68 Eir			Pb 207.19	Sn 118.69	32 Ge 72.59	Si 28.09	6 C 12.011	
	69 <b>Tm</b>			83 <b>Bi</b> 208.98	Sb 123.75	33 As 74.92	15 P 30.97	7 Z	
	70 <b>Yh</b> 173 04	٠		Po (209)	Te 127.60	34 Se 78.96	32.06	8 O	
_	71 Lui			85 <b>At</b> (210)	53 1 126.90		CI 35.45	9 F 19.00	
				86 <b>R.n</b> (222)	\$4 <b>X.e</b> 131.30	36 <b>I</b> (1) 83.80	18 <b>A1</b> : 39.95	10 <b>Ne</b> 20.18	2 <b>I·le</b> 4.003

\*Symbol (and name) provisional.

Actinides

9) **P**a

**Z** 

94 **Pu** 

95 **A**m

232.04

238.03

(244)

(247)

Numbers in parentheses: available radioactive isotope of longest half-life.

100 **Fin** (257)

101 **Md** (258)

No (259)

103 **L.j.** (260)